arXiv:1508.03922v4 [math.AG] 27 May 2016

OKOUNKOV BODIES ASSOCIATED TO PSEUDOEFFECTIVE DIVISORS

SUNG RAK CHOI, YOONSUK HYUN, JINHYUNG PARK, AND JOONYEONG WON

Abstract. An Okounkov body is a convex body in Euclidean space associated to a big divisor on a smooth projective variety with respect to an admissible flag. We introduce two different convex bodies associated to pseudoeffective divisors, called the valuative Okounkov bodies and limiting Okounkov bodies. As in the case with big divisors, these convex bodies reflect asymptotic properties of pseudoeffective divisors.

1. Introduction

For a divisor D on a smooth projective variety X of dimension n, one can associate a convex body $\Delta_{Y_\bullet}(D)$, called the Okounkov body of D, in the Euclidean space $\mathbb{R}^n$ with respect to an admissible flag $Y_\bullet$. After the pioneering works by Lazarsfeld-Mustaţă ([LM]) and Kaveh-Khovanskii ([KK]), motivated by earlier works by Okounkov ([O1], [O2]), the Okounkov bodies $\Delta_{Y_\bullet}(D)$ have received a considerable amount of attention in a variety of flavors. It is believed that this convex body carries rich information of the asymptotic invariants of D. For example, it was proven in [LM, Theorem A] that if D is big, then the Euclidean volume of $\Delta_{Y_\bullet}(D)$ in $\mathbb{R}^n$ is equal to the volume $\mathrm{vol}(D)$ of D up to the constant $n!$. However, little is known about the Okounkov bodies for non-big pseudoeffective divisors. One of the annoying phenomena is that for a pseudoeffective divisor D which is not big, the associated convex body $\Delta_{Y_\bullet}(D)$ is not full dimensional in $\mathbb{R}^n$ so that its Euclidean volume in $\mathbb{R}^n$ is zero. Nevertheless, it is still tempting to study asymptotic properties of pseudoeffective divisors using the associated Okounkov bodies.

In this paper, we introduce and study two different convex bodies, the valuative Okounkov body $\Delta^{\mathrm{val}}_{Y_\bullet}(D)$ and the limiting Okounkov body $\Delta^{\lim}_{Y_\bullet}(D)$, associated to a pseudoeffective divisor D. These convex bodies can be regarded as a generalization of the Okounkov body of a big divisor because if D is big, then we have

$$\Delta^{\mathrm{val}}_{Y_\bullet}(D) = \Delta^{\lim}_{Y_\bullet}(D) = \Delta_{Y_\bullet}(D).$$

We will readily see that $\Delta^{\mathrm{val}}_{Y_\bullet}(D)$ and $\Delta^{\lim}_{Y_\bullet}(D)$ with respect to a *suitable choice of an admissible flag* $Y_\bullet$ encode the asymptotic invariants of the divisor D as in the case of big divisors.

Turning to the details, we first recall the construction of the Okounkov body of a big divisor which is equivalent to the ones given by Lazarsfeld-Mustaţă ([LM]) and Kaveh-Khovanskii ([KK]). Let D be a big divisor on a smooth projective variety X of dimension n. Fix an admissible flag on X

$$Y_\bullet : \; X = Y_0 \supseteq Y_1 \supseteq \cdots \supseteq Y_{n-1} \supseteq Y_n = \{x\}$$

where each Y_i is an $(n-i)$-dimensional irreducible subvariety of X and it is nonsingular at x. For $D' \in |D|_{\mathbb{R}} := \{D' \mid D \sim_{\mathbb{R}} D' \geq 0\}$, we define a valuation-like function

$$\nu(D') = \nu_{Y_\bullet}(D') := \big(\nu_1(D'), \nu_2(D'), \cdots, \nu_n(D')\big) \in \mathbb{R}^n_{\geq 0}$$

as follows. Set $\nu_1(D') := \mathrm{ord}_{Y_1}(D'), \nu_2(D') := \mathrm{ord}_{Y_2}((D' - \nu_1(D')Y_1)|_{Y_1})$, and define ν_i for $i \geq 3$ inductively. The *Okounkov body* $\Delta_{Y_\bullet}(D)$ of D with respect to $Y_\bullet$ is defined as the closure of the convex hull of $\nu_{Y_\bullet}(|D|_{\mathbb{R}})$ in $\mathbb{R}^n_{\geq 0}$. Note that if a given divisor D satisfies $|D|_{\mathbb{R}} \neq \emptyset$ (or

---

Date: May 30, 2016.

Key words and phrases. Okounkov body, pseudoeffective divisor, Kodaira dimension, restricted volume, Nakayama subvariety, positive volume subvariety.

equivalently the Iitaka dimension $\kappa(D)$ is nonnegative), then the above construction can be carried out without any changes. See Subsection 3.1 for more details.

We now introduce the valuative Okounkov body of an effective divisor.

Definition 1.1. Let D be an $\mathbb{R}$-divisor with $\kappa(D) \geq 0$ on a smooth projective variety X, and $Y_\bullet$ be an admissible flag on X. The Okounkov body $\Delta_{Y_\bullet}(D)$ of D with respect to $Y_\bullet$ is called the *valuative Okounkov body* of D, and it is denoted by $\Delta^{\mathrm{val}}_{Y_\bullet}(D)$.

We note that $\Delta^{\mathrm{val}}_{Y_\bullet}(D)$ is not a numerical invariant (see Remark 3.13) but an invariant of linear equivalence (see Proposition 3.15). To extract asymptotic invariants of an effective divisor D from the valuative Okounkov body $\Delta^{\mathrm{val}}_{Y_\bullet}(D)$, we consider an admissible flag $Y_\bullet$ which contains a special subvariety $U \subseteq X$ such that $\kappa(D) = \dim U$ and the natural map $H^0(X, \mathcal{O}_X(\lfloor mD \rfloor)) \to H^0(U, \mathcal{O}_U(\lfloor mD|_U \rfloor))$ is injective for every integer $m \geq 0$. We call such U a *Nakayama subvariety* of D (see Subsection 2.3). We prove that any general subvariety of dimension $\kappa(D)$ is a Nakayama subvariety of D (Proposition 2.9). If D is big, then X itself is the unique Nakayama subvariety of D.

Theorem A (=Theorem 3.12). *Let D be an $\mathbb{R}$-divisor with $\kappa(D) \geq 0$ on a smooth projective variety X. Fix an admissible flag $Y_\bullet$ containing a Nakayama subvariety U of D such that $Y_n = \{x\}$ is a general point. Then we have*

$$\dim \Delta^{\mathrm{val}}_{Y_\bullet}(D) = \kappa(D) \text{ and } \mathrm{vol}_{\mathbb{R}^{\kappa(D)}}(\Delta^{\mathrm{val}}_{Y_\bullet}(D)) = \frac{1}{\kappa(D)!} \mathrm{vol}_{X|U}(D).$$

For convenience, we define dim(point) := 0 and $\mathrm{vol}_{\mathbb{R}^0}$(point) := 1 throughout the paper.

We associate another convex body $\Delta^{\lim}_{Y_\bullet}(D)$ to a pseudoeffective divisor D which reflect the asymptotic and numerical properties of D.

Definition 1.2. Let D be an $\mathbb{R}$-divisor with $\kappa_\nu(D) \geq 0$ on a smooth projective variety X, and $Y_\bullet$ be an admissible flag on X. The *limiting Okounkov body* of D with respect to $Y_\bullet$ is defined as

$$\Delta^{\lim}_{Y_\bullet}(D) := \lim_{\varepsilon \to 0+} \Delta_{Y_\bullet}(D + \varepsilon A) = \bigcap_{\varepsilon > 0} \Delta_{Y_\bullet}(D + \varepsilon A)$$

where A is an ample divisor on X.

Recall that D is a pseudoeffective divisor if and only if the numerical Iitaka dimension $\kappa_\nu(D)$ is nonnegative. To investigate the asymptotic properties of a pseudoeffective divisor D using the limiting Okounkov body $\Delta^{\lim}_{Y_\bullet}(D)$, we consider an admissible flag $Y_\bullet$ which contains a special subvariety $V \subseteq X$ such that $\kappa_\nu(D) = \dim V$, $V \not\subseteq \mathbf{B}_-(D)$ and $\mathrm{vol}^+_{X|V}(D) > 0$. We call such V a *positive volume subvariety* of D (see Subsection 2.4). We refer to Subsection 2.2 for the definition of the *augmented restricted volume* $\mathrm{vol}^+_{X|V}(D)$. We show that the intersection of $(n - \kappa_\nu(D))$ general ample divisors is a positive volume subvariety of D (Proposition 2.17). Thus we may take a *general* admissible flag $Y_\bullet$ constructed by successively intersecting general ample divisors. If D is big, then X itself is the unique positive volume subvariety of D.

Theorem B (=Corollary 3.22). *Let D be an $\mathbb{R}$-divisor with $\kappa_\nu(D) \geq 0$ on a smooth projective variety X. Fix an admissible flag $Y_\bullet$ containing a positive volume subvariety V of D. Then we have*

$$\dim \Delta^{\lim}_{Y_\bullet}(D) = \kappa_\nu(D) \quad \text{and} \quad \mathrm{vol}_{\mathbb{R}^{\kappa_\nu(D)}}(\Delta^{\lim}_{Y_\bullet}(D)) = \frac{1}{\kappa_\nu(D)!} \mathrm{vol}^+_{X|V}(D).$$

We can also extend the results of [LM, Proposition 4.1] and [J, Theorem A] to the pseudoeffective divisor case using the limiting Okounkov bodies. Note that the following theorem does not hold for the valuative Okounkov bodies (see Remarks 3.13 and 3.16).

Theorem C (=Theorem 3.28)**.** *Let D, D' be pseudoeffective $\mathbb{R}$-divisors on a smooth projective variety X. Then $D \equiv D'$ if and only if $\Delta^{\lim}_{Y_\bullet}(D) = \Delta^{\lim}_{Y_\bullet}(D')$ for every admissible flag $Y_\bullet$.*

We finally remark that Boucksom also defined and studied the limiting Okounkov body (the numerical Okounkov body in his terminology) in [B]. See Remark 3.24 for more details on his results.

This paper is organized as follows. We start in Section 2 by collecting relevant basic facts on asymptotic base locus, augmented restricted volume, Nakayama subvariety, and positive volume subvariety. Section 3 is the main part of this paper. We first recall the construction of the Okounkov body $\Delta_{Y_\bullet}(D)$ of a big divisor D in Subsection 3.1. We then introduce and study the valuative Okounkov body $\Delta^{\mathrm{val}}_{Y_\bullet}(D)$ and the limiting Okounkov body $\Delta^{\lim}_{Y_\bullet}(D)$ of a pseudoeffective divisor D in Subsections 3.2 and 3.3, respectively. In Section 4, we exhibit various examples.

Aknowledgement. We would like to thank Dr. A. M. Fulger for helpful suggestions and comments.

2. Preliminaries

In this section, we collect basic facts and introduce some notions used throughout the paper. First, we fix some notations. Throughout the paper, we work over the field $\mathbb{C}$ of complex numbers. For simplicity, a *variety* in this paper is assumed to be smooth, projective, reduced and irreducible, but a *subvariety* can be singular. By a *divisor* on a variety X, we always mean an $\mathbb{R}$-divisor unless otherwise stated. A divisor D is called *pseudoeffective* if its numerical class $[D] \in \mathrm{N}^1(X)_{\mathbb{R}} := \mathrm{N}^1(X) \otimes \mathbb{R}$ lies in $\overline{\mathrm{Eff}}(X)$, the closure of the cone $\mathrm{N}^1(X)_{\mathbb{R}}$ spanned by the classes of effective divisors. A divisor is called *big* if its numerical class lies in $\mathrm{Big}(X)$, the interior of $\overline{\mathrm{Eff}}(X)$.

2.1. **Asymptotic base locus.**

Let D be a pseudoeffective divisor on a variety X. When D is a $\mathbb{Q}$-divisor, we define the *stable base locus* of D as

$$\mathrm{SB}(D) := \bigcap_m \mathrm{Bs}(mD)$$

where the intersection is taken over all positive integers m such that mD are $\mathbb{Z}$-divisors, and $\mathrm{Bs}(mD)$ denotes the base locus of the linear system $|mD|$. The *augmented base locus* $\mathbf{B}_+(D)$ is defined as

$$\mathbf{B}_+(D) := \bigcap_A \mathrm{SB}(D - A)$$

where the intersection is taken over all ample divisors A such that $D - A$ are $\mathbb{Q}$-divisors. The *restricted base locus* $\mathbf{B}_-(D)$ of D is defined as

$$\mathbf{B}_-(D) := \bigcup_A \mathrm{SB}(D + A).$$

where the union is taken over all ample divisors A such that $D + A$ are $\mathbb{Q}$-divisors. We have $\mathbf{B}_-(D) \subseteq \mathrm{SB}(D) \subseteq \mathbf{B}_+(D)$ for a $\mathbb{Q}$-divisor D. One can check that a divisor D is ample (or nef) if and only if $\mathbf{B}_+(D) = \emptyset$ (respectively, $\mathbf{B}_-(D) = \emptyset$). The asymptotic base loci $\mathbf{B}_+(D)$ and $\mathbf{B}_-(D)$ depend only on the numerical class of D while $\mathrm{SB}(D)$ does not (see [La, Example 10.3.3]). For more properties, we refer to [ELMNP1].

Let $V \subseteq X$ be an irreducible subvariety of X. When D is big, we define *the asymptotic valuation* of V at D as

$$\mathrm{ord}_V(||D||) := \inf\{\mathrm{ord}_V(D') \mid D \equiv D' \geq 0\}.$$

When D is only pseudoeffective, we define $\mathrm{ord}_V(||D||) := \lim_{\varepsilon \to 0+} \mathrm{ord}_V(||D + \varepsilon A||)$ for some ample divisor A. This definition is independent of the choice of A, and the value $\sigma_{\mathrm{num}}(D)$

depends only on the numerical class $[D] \in \mathrm{N}^1(X)_{\mathbb{R}}$. It is a birational invariant: for a birational morphism $f : Y \to X$ with the exceptional divisor E such that $f(E) = V$, we have $\mathrm{ord}_V(||D||) = \mathrm{ord}_E(||f^*(D)||)$ (see [BBP, Lemma 1.4]). The restricted base locus $\mathbf{B}_-(D)$ can be characterized as follows.

Theorem 2.1 ([ELMNP1, Proposition 2.8],[N, Lemma V.1.9])**.** *Let D be a pseudoeffective divisor on a variety X, and $V \subseteq X$ an irreducible subvariety. Then $V \subseteq \mathbf{B}_-(D)$ if and only if* $\mathrm{ord}_V(||D||) > 0$.

For more details on asymptotic valuations, see [ELMNP1] and [N].

2.2. **Augmented restricted volume.**

To introduce the notion of the augmented restricted volume, we first recall the classical volume function and the restricted volume function. Let D be a $\mathbb{Q}$-divisor on a variety X of dimension n. Recall that the *volume* of D is defined as

$$\mathrm{vol}_X(D) := \limsup_{m\to\infty} \frac{h^0(X, \mathcal{O}_X(mD))}{m^n/n!}.$$

The volume $\mathrm{vol}_X(D)$ depends only on the numerical class of D. Furthermore, this function uniquely extends to a continuous function

$$\mathrm{vol}_X : \mathrm{Big}(X) \to \mathbb{R}.$$

Let V be a v-dimensional subvariety of X. For the natural restriction map $\varphi : H^0(X, \mathcal{O}_X(D)) \to H^0(V, \mathcal{O}_V(D))$, we denote the image $\mathrm{Im}(\varphi)$ by $H^0(X|V, D)$ and its dimension by $h^0(X|V, D)$. The *restricted volume* of D along V is defined as

$$\mathrm{vol}_{X|V}(D) := \limsup_{m\to\infty} \frac{h^0(X|V, mD)}{m^v/v!}$$

([ELMNP2, Definition 2.1]). The restricted volume $\mathrm{vol}_{X|V}(D)$ depends only on the numerical class of D. Furthermore, this function uniquely extends to a continuous function

$$\mathrm{vol}_{X|V} : \mathrm{Big}^V(X) \to \mathbb{R}$$

where $\mathrm{Big}^V(X)$ is the set of all $\mathbb{R}$-divisor classes ξ such that V is not properly contained in any irreducible component of $\mathbf{B}_+(\xi)$. Note that by letting $V = X$, we recover the usual volume function $\mathrm{vol}_{X|X}(D) = \mathrm{vol}_X(D)$. For more details on the functions vol_X and $\mathrm{vol}_{X|V}$, see [ELMNP2], [La], [Le], etc.

By [ELMNP2, Theorem 5.2], $\mathrm{vol}_{X|V}(D) = 0$ if V is an irreducible component of $\mathbf{B}_+(D)$. Similarly, $\mathrm{vol}_X(D) = 0$ if D is not big (i.e., $X = \mathbf{B}_+(D)$). Thus the functions $\mathrm{vol}_{X|V}$ and vol_X do not capture the subtle asymptotic properties of pseudoeffective divisors that are not big on V or X. In such situations, the following function seems useful.

Definition 2.2. Let D be a pseudoeffective divisor on X, and $V \subseteq X$ be a subvariety such that $V \not\subseteq \mathbf{B}_-(D)$. The *augmented restricted volume* of D along V is defined as

$$\mathrm{vol}^+_{X|V}(D) := \lim_{\varepsilon\to 0+} \mathrm{vol}_{X|V}(D + \varepsilon A)$$

where A is an ample divisor on X.

The definition is independent of the choice of A. As with vol_X and $\mathrm{vol}_{X|V}$, one can check that the augmented restricted volume $\mathrm{vol}^+_{X|V}(D)$ depends only on the numerical class of D. By the continuity of the function $\mathrm{vol}_{X|V}$, we see that $\mathrm{vol}^+_{X|V}(D)$ coincides with $\mathrm{vol}_{X|V}(D)$ for $D \in \mathrm{Big}^V(X)$. Furthermore, we have a continuous function

$$\mathrm{vol}^+_{X|V} : \overline{\mathrm{Eff}}^V(X) \to \mathbb{R}$$

where $\overline{\mathrm{Eff}}^V(X) := \mathrm{Big}^V(X) \cup \{\xi \in \overline{\mathrm{Eff}}(X) \setminus \mathrm{Big}(X) \mid V \not\subseteq \mathbf{B}_-(\xi)\}$. For $D \in \overline{\mathrm{Eff}}^V(X)$, the following inequalities hold by definition:

$$\mathrm{vol}_{X|V}(D) \leq \mathrm{vol}^+_{X|V}(D) \leq \mathrm{vol}_V(D|_V).$$

Both inequalities can be strict in general.

Example 2.3. Let S be a relatively minimal rational elliptic surface, and H be an ample divisor on S. Take a general element $V \in |kH|$ for a sufficiently large $k > 0$. Then $\mathrm{vol}_{S|V}(-K_S) = \limsup\limits_{m\to\infty} \frac{h^0(S,\mathcal{O}_S(-mK_S))}{m}$ which is independent of k and H. However, one can see that $\mathrm{vol}^+_{S|V}(-K_S) = k(-K_S)\cdot H$ which can be arbitrarily large depending on k. Thus $\mathrm{vol}_{S|V}(-K_S) < \mathrm{vol}^+_{S|V}(-K_S)$.

2.3. **Nakayama subvariety.**

In this subsection, we introduce and study Nakayama subvarieties of divisors, which are closely related to the Iitaka dimension.

Definition 2.4. For a divisor D on a variety X, let $\mathbb{N}(D) = \{m \in \mathbb{Z}_{>0} | \, |\lfloor mD \rfloor| \neq \emptyset\}$. For $m \in \mathbb{N}(D)$, let $\Phi_{mD} : X \dashrightarrow \mathbb{P}^{\dim|\lfloor mD\rfloor|}$ be the rational map defined by the linear system $|\lfloor mD \rfloor|$. We define the *Iitaka dimension* of D as the following value

$$\kappa(D) := \begin{cases} \max\{\dim \mathrm{Im}(\Phi_{mD}) \mid m \in \mathbb{N}(D)\} & \text{if } \mathbb{N}(D) \neq \emptyset \\ -\infty & \text{if } \mathbb{N}(D) = \emptyset. \end{cases}$$

Remark 2.5. If D is a divisor such that $\kappa(D) \geq 0$, then there exists an integer $m_0 \geq 1$ such that $h^0(X, \mathcal{O}_X(\lfloor mm_0D \rfloor)) \sim m^{\kappa(D)}$ for $m \gg 0$ ([N, Theorem II.3.7]). For this m_0, we have $\kappa(D) = \kappa(m_0D)$. Thus in many situations below, we may assume that $m_0 = 1$.

Remark 2.6. If $D \sim_{\mathbb{R}} D'$ and D, D' are effective, then $\kappa(D) = \kappa(D')$. However, the Iitaka dimension $\kappa(D)$ is neither an $\mathbb{R}$-linear nor numerical invariant in general ([G, Remark 2.1], [Le, Example 6.1]).

Definition 2.7. Let D be a divisor on a variety X such that $\kappa(D) \geq 0$. A subvariety $U \subseteq X$ is called a *Nakayama subvariety* of D if $\kappa(D) = \dim U$ and the natural map

$$H^0(X, \mathcal{O}_X(\lfloor mD \rfloor)) \to H^0(U, \mathcal{O}_U(\lfloor mD|_U \rfloor))$$

is injective (or equivalently, $H^0(X, \mathcal{I}_U \otimes \mathcal{O}_X(\lfloor mD \rfloor)) = 0$ where $\mathcal{I}_U$ is an ideal sheaf of U in X) for every integer $m \geq 0$.

By definition, $U = X$ is the (unique) Nakayama subvariety of D if and only if D is big. More generally, Nakayama subvarieties have the following property.

Lemma 2.8. *Let D be a divisor such that $\kappa(D) \geq 0$, and $U \subseteq X$ be its Nakayama subvariety. Then $D|_U$ is big.*

Proof. By the definition of Nakayama subvariety, we have

$$h^0(U, \mathcal{O}_U(\lfloor mD|_U \rfloor)) \geq h^0(X, \mathcal{O}_X(\lfloor mD \rfloor))$$

for every integer $m \geq 0$. Since $h^0(X, \mathcal{O}_X(\lfloor mD \rfloor)) \sim m^{\kappa(D)}$ for $m \gg 0$, the assertion follows. ☐

The following lemma guarantees the existence of Nakayama subvarieties. It also shows that a Nakayama subvariety is not unique in general.

Proposition 2.9 (cf. [N, Lemma V.2.11])**.** *Let D be a divisor on X such that $\kappa(D) \geq 0$. Any general subvariety $U \subseteq X$ of dimension $\kappa(D)$ is a Nakayama subvariety of D.*

Proof. Consider a dominant rational map $\varphi_m : X \dashrightarrow Z_m$ induced by a complete linear series $|\lfloor mD \rfloor|$ for any integer $m > 0$ such that $|\lfloor mD \rfloor| \neq \emptyset$. We have $\dim Z_m \leq \kappa(D) = \dim U$. Let $f : Y \to X$ be the blow-up at U with the exceptional divisor E. Since $U \subseteq X$ is general, $f^*(\lfloor mD \rfloor) - kE$ is not effective for any $k > 0$. Thus $H^0(X, \mathcal{I}_U \otimes \mathcal{O}_X(\lfloor mD \rfloor)) = 0$, and hence, the assertion follows. □

Corollary 2.10. *If the effective divisors D, D' on a variety X satisfy $D \sim_{\mathbb{R}} D'$, then D, D' have a common Nakayama subvariety.*

Proof. The given condition implies that $\kappa(D) = \kappa(D') \geq 0$. By proposition 2.9, any general subvariety $U \subseteq X$ of dimension $\kappa(D)$ is a Nakayama subvariety of D. □

2.4. **Positive volume subvariety.**

We introduce and study positive volume subvarieties of divisors, which are closely related to the numerical Iitaka dimension and the restricted volume. First, we review the numerical Iitaka dimension.

Definition 2.11. Let D be a divisor on a variety X. We define the *numerical Iitaka dimension* of D as the nonnegative integer

$$\kappa_\nu(D) := \max\left\{ k \in \mathbb{Z}_{\geq 0} \,\middle|\, \limsup_{m\to\infty} \frac{h^0(X, \mathcal{O}_X(\lfloor mD \rfloor + A))}{m^k} > 0 \right\}$$

for some fixed ample Cartier divisor A if $h^0(X, \mathcal{O}_X(\lfloor mD \rfloor + A)) \neq \emptyset$ for infinitely many $m > 0$ and we let $\kappa_\nu(D) := -\infty$ otherwise.

The numerical Iitaka dimension $\kappa_\nu(D)$ depends only on the numerical class $[D] \in \mathrm{N}^1(X)_{\mathbb{R}}$. One can easily check that $\kappa(D) \leq \kappa_\nu(D)$ holds and the inequality is strict in general (see [Le, Example 6.1]). However, if $\kappa_\nu(D) = \dim X$, then $\kappa(D) = \dim X$. See [Le] and [N] for detailed properties of κ and κ_ν.

The following theorem relates the numerical Iitaka dimension $\kappa_\nu(D)$ and $\mathrm{vol}^+_{X|V}(D)$.

Theorem 2.12 ([Le, Theorem 1.1]). *Let D be a pseudoeffective divisor on a variety X. Then we have*

$$\kappa_\nu(D) = \max\{\dim W |\ \mathrm{vol}^+_{X|W}(D) > 0 \text{ and } W \not\subseteq \mathbf{B}_-(D)\}.$$

The numerical Iitaka dimension $\kappa_\nu(D)$ actually coincides with many other invariants defined with D. For details, see [Le].

Definition 2.13. Let D be a pseudoeffective divisor on a variety X. A subvariety $V \subseteq X$ of dimension $\kappa_\nu(D)$ such that $\mathrm{vol}^+_{X|V}(D) > 0$ and $V \not\subseteq \mathbf{B}_-(D)$ is called a *positive volume subvariety* of D. (In other words, a positive volume subvariety $V \subseteq X$ is the subvariety at which the maximum in Theorem 2.12 is attained.)

Theorem 2.12 guarantees the existence of a positive volume subvariety.

Example 2.14. Let S be a relatively minimal rational surface with a reducible singular fiber, and E a (-2)-curve in a singular fiber. Then $E \not\subseteq \mathbf{B}_-(-K_S)$ but $\mathrm{vol}^+_{S|E}(-K_S) = 0$. In this example, we see that $V \not\subseteq \mathbf{B}_-(D)$ and $\dim V = \kappa_\nu(D)$ does not imply $\mathrm{vol}^+_{X|V}(D) > 0$.

By definition, if D is a big divisor on X, then $V = X$ is the unique positive volume subvariety of D. We prove some notable properties of positive volume subvariety.

Lemma 2.15. *If D and D' are pseudoeffective divisors on a variety X such that $D \equiv D'$, then they have the same positive volume subvarieties.*

Proof. Note first that $\mathbf{B}_-(D) = \mathbf{B}_-(D')$. The statement then follows from the fact that $\mathrm{vol}^+_{X|V}(D)$ depends only on the numerical class $[D]$ for any $V \not\subseteq \mathbf{B}_-(D) = \mathbf{B}_-(D')$. □

Let V be a smooth subvariety of a variety X. A birational morphism $f : Y \to X$ is said to be *V-birational* if V is not contained in the center of the f-exceptional locus. We denote by $\widetilde{V}$ the proper transform of V on Y. Then the pair $(Y, \widetilde{V})$ is called a *V-birational model* of X (see [Le, Definition 2.10]).

Proposition 2.16. *Let $V \subseteq X$ be a positive volume subvariety of a pseudoeffective divisor D on a variety X. If $f : Y \to X$ is a V-birational morphism, then $\widetilde{V}$ is also a positive volume subvariety of f^*D.*

Proof. The birational transform $\widetilde{V} \subseteq Y$ of V has dimension $\dim \widetilde{V} = \dim V = \kappa_\nu(D) = \kappa_\nu(f^*D)$. If $\widetilde{V} \subseteq \mathbf{B}_-(f^*(D))$, then $V = f(\widetilde{V}) \subseteq f(\mathbf{B}_-(f^*(D)) = \mathbf{B}_-(D)$. Thus $\widetilde{V} \not\subseteq \mathbf{B}_-(f^*(D))$. It is enough to show that $\operatorname{vol}^+_{Y|\widetilde{V}}(f^*D) > 0$. Since $V \not\subseteq \mathbf{B}_-(D)$ by definition, for any ample divisor A on X, we have $V \not\subseteq \mathbf{B}_+(D + A)$. We can also check that $\widetilde{V} \not\subseteq \mathbf{B}_+(f^*D + f^*A)$ using [BBP, Proposition 2.3]. Note that there is an effective divisor E on Y such that $f^*A - E$ is ample. Thus we have

$$\operatorname{vol}^+_{Y|\widetilde{V}}(f^*D) = \lim_{\varepsilon \to 0+} \operatorname{vol}_{Y|\widetilde{V}}(f^*D + \varepsilon(f^*A - E)).$$

The difference $|\operatorname{vol}_{Y|\widetilde{V}}(f^*D + \varepsilon(f^*A - E)) - \operatorname{vol}_{Y|\widetilde{V}}(f^*D + \varepsilon f^*A)|$ can be arbitrarily small when $\varepsilon > 0$ is sufficiently small. On the other hand, by [ELMNP2, Lemma 2.4], we have

$$\operatorname{vol}^+_{X|V}(D) = \lim_{\varepsilon \to 0+} \operatorname{vol}_{X|V}(D + \varepsilon A) = \lim_{\varepsilon \to 0+} \operatorname{vol}_{Y|\widetilde{V}}(f^*D + \varepsilon f^*A).$$

Since $\operatorname{vol}^+_{X|V}(D) > 0$, it follows that $\operatorname{vol}^+_{Y|\widetilde{V}}(f^*D) > 0$. □

The following proposition guarantees the existence of positive volume subvarieties.

Proposition 2.17. *Let D be a pseudoeffective divisor on a variety X of dimension n. If $\kappa_\nu(D) < n$, then the subvariety $V := A_1 \cap \cdots \cap A_{n-\kappa_\nu(D)}$ satisfies $\operatorname{vol}^+_{X|V}(D) > 0$, where A_i are sufficiently ample divisors. In particular, V is a positive volume subvariety of D.*

Proof. By Theorem 2.12, there exists a positive volume subvariety W of D. In particular, we have $\operatorname{vol}^+_{X|W}(D) > 0$. We can take a sequence $\{H_i\}_{i \in \mathbb{Z}_{\geq 0}}$ of ample divisors on X such that each $D + H_i$ is a $\mathbb{Q}$-divisor and $H_i \to 0$ as $i \to \infty$. For a large and sufficiently divisible integer k, choose $\kappa_\nu(D)$ general divisors $E^1_{k,i}, \cdots, E^{\kappa_\nu(D)}_{k,i} \in |k(D + H_i)|$. We have

$$\#(V \cap E^1_{k,i} \cap \cdots \cap E^{\kappa_\nu(D)}_{k,i} \setminus \operatorname{SB}(D + H_i)) \geq \#(W \cap E^1_{k,i} \cap \cdots \cap E^{\kappa_\nu(D)}_{k,i} \setminus \operatorname{SB}(D + H_i)).$$

Thus by applying [ELMNP2, Theorem B], we obtain

$$\begin{aligned} \operatorname{vol}_{X|V}(D + H_i) &= \lim_{k \to \infty} \frac{\#(V \cap E^1_{k,i} \cap \cdots \cap E^{\kappa_\nu(D)}_{k,i} \setminus \operatorname{SB}(D+H_i))}{k^{\kappa_\nu(D)}} \\ &\geq \lim_{k \to \infty} \frac{\#(W \cap E^1_{k,i} \cap \cdots \cap E^{\kappa_\nu(D)}_{k,i} \setminus \operatorname{SB}(D+H_i))}{k^{\kappa_\nu(D)}} = \operatorname{vol}_{X|W}(D + H_i). \end{aligned}$$

This immediately implies that $\operatorname{vol}^+_{X|V}(D) \geq \operatorname{vol}^+_{X|W}(D) > 0$. Since $A_1, \cdots, A_{n-\kappa_\nu(D)}$ are sufficiently ample, we can assume that $V \not\subseteq \mathbf{B}_-(D)$. □

It is known that $D|_V$ is pseudoeffective for any subvariety $V \subseteq X$ such that $V \not\subseteq \mathbf{B}_-(D)$. Thus $\kappa(D|_V) \geq 0$. Furthermore, we have the following.

Theorem 2.18. *Let V be a positive volume subvariety of a pseudoeffective divisor D on a variety X. Then $\operatorname{vol}_V(D|_V) > 0$. In particular, $D|_V$ is big on V and $\kappa_\nu(D) = \kappa_\nu(D|_V) = \dim V$.*

Proof. By taking a suitable V-birational model and considering Proposition 2.16, we may assume that V is smooth. We can take a sequence of ample divisors A_i such that $D+A_i$ are $\mathbb{Q}$-divisors and $A_i \to 0$ as $i \to \infty$. Since $V \subseteq X$ is a positive volume subvariety of D, there exists a constant C_0 such that for any large integer $k > 0$, we have

$$\limsup_{m\to\infty} \frac{h^0(X|V, m(D+A_k))}{m^{\kappa_\nu(D)}} > C_0$$

where m is taken over all positive integers such that $m(D+A_k)$ is a $\mathbb{Z}$-divisor. Note that $h^0(X|V, m(D+A_k)) \leq h^0(V, m(D|_V + A_k|_V))$. Thus for any large integer $k > 0$, we also have

$$\limsup_{m\to\infty} \frac{h^0(V, m(D|_V + A_k|_V))}{m^{\kappa_\nu(D)}} > C_0.$$

By the continuity of the volume function vol_V, we get $\mathrm{vol}_V(D|_V) > 0$. Thus $D|_V$ is big on V. □

3. Okounkov body of a pseudoeffective divisor

In this section, we introduce and study two Okounkov bodies $\Delta^{\mathrm{val}}_{Y_\bullet}(D)$ and $\Delta^{\mathrm{lim}}_{Y_\bullet}(D)$ for a pseudoeffective divisor D on a smooth projective variety of dimension n.

3.1. Okounkov body $\Delta_{Y_\bullet}(D)$.

We first recall the construction of the Okounkov body of a big divisor D and a graded linear series $W_\bullet$ associated to D from [LM]. We remark that the construction of $\Delta_{Y_\bullet}(D)$ in the introduction is equivalent to the one given below (see [KL]).

Throughout this subsection, we fix an *admissible flag* $Y_\bullet$ on a variety X of dimension n

$$Y_\bullet : \ X = Y_0 \supset Y_1 \supset Y_2 \supset \cdots \supset Y_{n-1} \supset Y_n = \{x\}$$

where each Y_i is an $(n-i)$-dimensional subvariety of X and it is nonsingular at x. We first assume that D is a big Cartier divisor on X. For a section $s \in H^0(X, \mathcal{O}_X(D)) \setminus \{0\}$, we define the function

$$\nu(s) = \nu_{Y_\bullet}(s) := (\nu_1(s), \nu_2(s), \cdots, \nu_n(s)) \in \mathbb{Z}^n_{\geq 0}$$

as follows. Let $\nu_1 = \nu_1(s) := \mathrm{ord}_{Y_1}(s)$. Using a local equation f_1 for Y_1 in X, define a section

$$s_1' = s \otimes f_1^{-\nu_1} \in H^0(X, \mathcal{O}_X(D - \nu_1 Y_1)).$$

Since s_1' does not vanish identically along Y_1, its restriction $s_1'|_{Y_1}$ defines a nonzero section

$$s_1 := s_1'|_{Y_1} \in H^0(Y_1, \mathcal{O}_{Y_1}(D - \nu_1 Y_1)) \setminus \{0\}.$$

Now take $\nu_2 = \nu_2(s) := \mathrm{ord}_{Y_2}(s_1)$. Continuing this manner, we obtain the nonnegative integers $\nu_i(s)$ for all $1 \leq i \leq n$ so that $\nu(s) \in \mathbb{Z}^n_{\geq 0}$. Each integer $\nu_i = \nu_i(s)$ does not depend on the choice of the local equation f_i chosen to define s_i' and s_i.

Recall that a *graded linear series* $W_\bullet(D) = \{W_m := W_m(D)\}_{m\geq 0}$ associated to D consists of subspaces $W_m \subseteq H^0(X, \mathcal{O}_X(mD))$ with $W_0 = \mathbb{C}$ satisfying $W_m \cdot W_l \subseteq W_{m+l}$ for all $m, l \geq 0$. The *graded semigroup* of $W_\bullet$ is

$$\Gamma(W_\bullet) := \{(\nu(s), m) \mid \ W_m \neq \{0\} \text{ and } s \in W_m \setminus \{0\} \ \} \subseteq \mathbb{Z}^n_{\geq 0} \times \mathbb{Z}_{\geq 0}.$$

Definition 3.1. Under the same notations as above, we associate the convex body $\Delta_{Y_\bullet}(W_\bullet)$ called the *Okounkov body* of a graded linear series $W_\bullet$ with respect to an admissible flag $Y_\bullet$ on X as follows:

$$\Delta_{Y_\bullet}(W_\bullet) := \Sigma(\Gamma(W_\bullet)) \cap (\mathbb{R}^n_{\geq 0} \times \{1\})$$

where $\Sigma(\Gamma(W_\bullet))$ denotes the closure of the convex cone in $\mathbb{R}^n_{\geq 0} \times \mathbb{R}_{\geq 0}$ spanned by $\Gamma(W_\bullet)$. If $W_\bullet$ is a complete graded linear series (i.e., $W_m = H^0(X, \mathcal{O}(mD))$ for each m), then we denote by $\Delta_{Y_\bullet}(D) = \Delta_{Y_\bullet}(W_\bullet)$.

We naturally treat $\Delta_{Y_\bullet}(W_\bullet)$ as a convex subset in $\mathbb{R}^n$, i.e., $\Delta_{Y_\bullet}(W_\bullet) \subseteq \mathbb{R}^n$. By construction, the Okounkov body $\Delta_{Y_\bullet}(D)$ depends on the choice of the admissible flag $Y_\bullet$. By the homogeneity of $\Delta_{Y_\bullet}(D)$ ([LM, Proposition 4.1]), we can extend the construction of $\Delta_{Y_\bullet}(D)$ to $\mathbb{Q}$-divisors D and even to $\mathbb{R}$-divisors using the continuity of $\Delta_{Y_\bullet}(D)$.

Remark 3.2. Note that the above constructions of $\Delta_{Y_\bullet}(D)$ and $\Delta_{Y_\bullet}(W_\bullet)$ work as long as $H^0(X, \mathcal{O}_X(mD)) \neq 0$ or $W_m \neq 0$ for some m. Thus it is not necessary to assume that a given divisor D is big to define the Okounkov body $\Delta_{Y_\bullet}(D)$.

Remark 3.3. Let $Y_\bullet$ be an admissible flag on a variety X of dimension n. For each integer k such that $0 \leq k \leq n$, we can define the k-th *partial flag* $Y_{k\bullet}$ of $Y_\bullet$ as

$$Y_{k\bullet} : Y_k \supseteq Y_{k+1} \supseteq \cdots \supseteq Y_n = \{x\}.$$

Then $Y_{k\bullet}$ is an admissible flag on a smooth projective variety Y_k of dimension $n-k$. Thus for a Cartier divisor D on X, we can define the function

$$\nu_{Y_{k\bullet}} : H^0(X|Y_k, \mathcal{O}_X(D)) \setminus \{0\} \to \mathbb{Z}^{n-k}_{\geq 0}$$

as we did above. Let

$$\Gamma_{Y_k}(D) := \{(\nu_{Y_{k\bullet}}(s), m) | s \in H^0(X|Y_k, \mathcal{O}_X(mD)) \setminus \{0\}, m \geq 0\}.$$

The *k-th partial Okounkov body* of a divisor D on X with respect to the k-th partial flag $Y_{k\bullet}$ is defined as

$$\Delta_{Y_{k\bullet}}(D) := \Sigma(\Gamma_{Y_k}(D)) \cap (\mathbb{R}^{n-k}_{\geq 0} \times \{1\})$$

where $\Sigma(\Gamma_{Y_k}(D))$ is the closure of the convex cone in $\mathbb{R}^{n-k}_{\geq 0} \times \mathbb{R}_{\geq 0}$ spanned by $\Gamma_{Y_k}(D)$. These partial flags and partial Okounkov bodies will be useful in the next subsections.

Under the following conditions, the Okounkov body $\Delta_{Y_\bullet}(W_\bullet)$ behaves well.

Definition 3.4 ([LM, Definitions 2.5 and 2.9])**.**

(1) We say that a graded linear series $W_\bullet$ satisfies *Condition* (B) if $W_m \neq 0$ for all $m \gg 0$, and for all sufficiently large m, the rational map $\varphi_m : X \dashrightarrow \mathbb{P}(W_m)$ defined by $|W_m|$ is birational onto its image.

(2) We say that a graded linear series $W_\bullet$ satisfies *Condition* (C) if
  i. for every $m \gg 0$, there exists an effective divisor F_m such that the divisor $A_m := mD - F_m$ is ample, and
  ii. for all sufficiently large p, we have

$$H^0(X, \mathcal{O}_X(pA_m)) \subseteq W_{pm} \subseteq H^0(X, \mathcal{O}_X(pmD)).$$

If $W_\bullet$ is complete, i.e., $W_m = H^0(X, \mathcal{O}_X(mD))$ for all $m \geq 0$ and D is big, then it automatically satisfies Condition (C).

Theorem 3.5 ([LM, Theorem 2.13])**.** *Suppose that a graded linear series $W_\bullet$ satisfies Condition (B) or Condition (C). Then for any admissible flag $Y_\bullet$ (with a general choice of the point $Y_n = \{x\}$ when $W_\bullet$ only satisfies Condition (B)), we have*

$$\dim \Delta_{Y_\bullet}(W_\bullet) = \dim X (= n) \quad \textit{and} \quad \mathrm{vol}_{\mathbb{R}^n}(\Delta_{Y_\bullet}(W_\bullet)) = \frac{1}{n!} \cdot \mathrm{vol}_X(W_\bullet)$$

where

$$\mathrm{vol}_X(W_\bullet) := \lim_{m \to \infty} \frac{\dim W_m}{m^n/n!}.$$

Note that [LM, Theorem 2.13] also requires Condition (A) ([LM, Definition 2.4]), but it is automatically satisfied since we always assume our variety X is projective. The smoothness assumption on X is not necessary for this theorem.

Remark 3.6. It is well known by [LM, Proposition 4.1] that for a fixed admissible flag $Y_\bullet$ on X, if D is a big divisor on X, then $\Delta_{Y_\bullet}(D)$ depends only on the numerical class of D. If D is not big, then it is no longer true. See Remark 3.13.

Remark 3.7. The k-th partial Okounkov body $\Delta_{Y_{k\bullet}}(D)$ depends on the numerical class of D if $D|_{Y_k}$ is big.

3.2. **Valuative Okounkov body $\Delta^{\text{val}}_{Y_\bullet}(D)$.**

Now we define the valuative Okounkov body $\Delta^{\text{val}}_{Y_\bullet}(D)$ of a divisor D with $\kappa(D) \geq 0$ on a variety X of dimension n.

Definition 3.8. Let D be a divisor on a variety X of dimension n such that $\kappa(D) \geq 0$. The *valuative Okounkov body* $\Delta^{\text{val}}_{Y_\bullet}(D)$ associated to D with respect to an admissible flag $Y_\bullet$ on X is defined as

$$\Delta^{\text{val}}_{Y_\bullet}(D) := \Delta_{Y_\bullet}(D) \subseteq \mathbb{R}^n.$$

For a divisor D with $\kappa(D) = -\infty$, we define $\Delta^{\text{val}}_{Y_\bullet}(D) := \emptyset$.

Obviously, the definition of $\Delta^{\text{val}}_{Y_\bullet}(D)$ depends on the choice of $Y_\bullet$. However, we will see below that if $Y_\bullet$ contains a Nakayama subvariety U of D (i.e., $Y_{n-\kappa(D)} = U$), the valuative Okounkov body $\Delta^{\text{val}}_{Y_\bullet}(D)$ depends only on the choice of a Nakayama subvariety of D and the $(n-\kappa(D))$-th partial flag on it.

Remark 3.9. If D is big, then $\Delta^{\text{val}}_{Y_\bullet}(D)$ coincides with $\Delta_{Y_\bullet}(D)$ for any admissible flag $Y_\bullet$ on X.

Now fix a Nakayama subvariety $U \subseteq X$ of D. Consider the *restricted complete graded linear series* $W_\bullet(D|U)$ of a Cartier divisor D along U which is given by $W_m(D|U) := H^0(X|U, mD)$ for each $m \geq 0$. Recall that $H^0(X|U, mD)$ is the image of the restriction map $\varphi_U : H^0(X, \mathcal{O}_X(mD)) \to H^0(U, \mathcal{O}_U(mD|_U))$. Note that $W_\bullet(D|U)$ is a graded linear subseries of $W_\bullet(D|_U)$ on U. Thus we can consider the function

$$\nu_{Y_{n-\kappa(D)\bullet}} : W_m(D|U) \setminus \{0\} \to \mathbb{Z}^{\kappa(D)}.$$

Since the natural restriction map φ_U is injective for each m, we may treat the function $\nu_{Y_{n-\kappa(D)\bullet}}$ as defined on the sections $s \in H^0(X, \mathcal{O}_X(mD)) \setminus \{0\}$ by letting

$$\nu_{Y_{n-\kappa(D)\bullet}}(s) := \nu_{Y_{n-\kappa(D)\bullet}}(\varphi_U(s)) \in \mathbb{Z}^{\kappa(D)}.$$

Proposition 3.10. *Let $s \in H^0(X, \mathcal{O}_X(mD)) \setminus \{0\}$. Then under the settings as above, we have*

$$\nu_{Y_\bullet}(s) = (\underbrace{0, \ldots, 0}_{n-\kappa(D)}, \nu_{Y_{n-\kappa(D)\bullet}}(s)).$$

In particular, $\Delta^{\text{val}}_{Y_\bullet}(D) \subseteq \{0\}^{n-\kappa(D)} \times \mathbb{R}^{\kappa(D)} \subseteq \mathbb{R}^n$ and we can regard $\Delta^{\text{val}}_{Y_\bullet}(D)$ as a subset of $\mathbb{R}^{\kappa(D)}$.

Proof. By the definition of Nakayama subvariety, we have for all $m \geq 0$

$$H^0(X, \mathcal{I}_U \otimes \mathcal{O}_X(mD)) = 0.$$

Let k be any integer such that $1 \leq k \leq n-\kappa(D)$. Then since $Y_k \supseteq U$, it follows that $H^0(X, \mathcal{I}_{Y_k} \otimes \mathcal{O}_X(mD)) = 0$. Thus we obtain $\nu_k(s) = 0$. This implies that

$$\nu_{n-\kappa(D)+1}(s) = \nu_{Y_{n-\kappa(D)\bullet},1}(s|_{Y_{n-\kappa(D)}}) = \nu_{Y_{n-\kappa(D)\bullet},1}(\varphi_U(s)).$$

This shows the required equality. □

Remark 3.11. Let $Y_\bullet$ be an admissible flag containing a Nakayama subvariety U of a pseudoeffective divisor D on X. Then the valuative Okounkov body $\Delta^{\text{val}}_{Y_\bullet}(D)$ can be constructed alternatively as follows. Consider the $(n-\kappa(D))$-th partial flag $Y_{n-\kappa(D)\bullet}$ on U. Since $W_\bullet(D|U)$ is a graded linear subseries associate to $D|_U$ on U, we can define the Okounkov body

$$\Delta_{Y_{n-\kappa(D)\bullet}}(W_\bullet(D|U)) \subseteq \mathbb{R}^{\kappa(D)}$$

as in Definition 3.1. By regarding $\Delta^{\text{val}}_{Y_\bullet}(D)$ as a subset of $\mathbb{R}^{\kappa(D)}$ (Proposition 3.10), we have

$$\Delta^{\text{val}}_{Y_\bullet}(D) = \Delta_{Y_{n-\kappa(D)\bullet}}(W_\bullet(D|U)).$$

Thus $\Delta_{Y_{n-\kappa(D)\bullet}}(W_\bullet(D|U))$ gives an alternative construction of $\Delta^{\text{val}}_{Y_\bullet}(D)$.

The following is the main property of $\Delta^{\text{val}}_{Y_\bullet}(D)$.

Theorem 3.12. *Let D be a divisor such that $\kappa(D) \geq 0$ on a smooth projective variety X, and fix an admissible flag $Y_\bullet$ containing a Nakayama subvariety U of D such that $Y_n = \{x\}$ is a general point. Then we have*

$$\dim \Delta^{\text{val}}_{Y_\bullet}(D) = \kappa(D) \quad \textit{and} \quad \text{vol}_{\mathbb{R}^{\kappa(D)}}(\Delta^{\text{val}}_{Y_\bullet}(D)) = \frac{1}{\kappa(D)!}\text{vol}_{X|U}(D).$$

Proof. We first consider the case where D is a Cartier divisor. We see that $\Delta^{\text{val}}_{Y_\bullet}(mD) = \frac{1}{m}\Delta^{\text{val}}_{Y_\bullet}(D)$ for any integer $m > 0$. Thus we may assume that the constant m_0 in Remark 2.5 is equal to 1. In particular, $h^0(X, \mathcal{O}_X(mD)) > 0$ for any integer $m > 0$. By Remark 3.11, we have $\Delta^{\text{val}}_{Y_\bullet}(D) = \Delta_{Y_{n-\kappa(D)\bullet}}(W_\bullet(D|U))$. By the properties of a Nakayama subvariety, there exits an integer m_0 such that $h^0(X, \mathcal{O}_X(mm_0D)) = \dim W_{mm_0} \sim m^{\kappa(D)}$ for all $m \gg 0$ (cf. Remark 2.5) and $\dim U = \kappa(D)$. Furthermore, $Y_n = \{x\}$ is assumed to be general. By [LM, Remark 2.8] and Theorem 3.5, we obtain the theorem. By the similar argument as in [LM], we can easily extend the results to $\mathbb{Q}$- or $\mathbb{R}$-divisors. We leave the details to the interested readers. □

Remark 3.13. It is natural to ask whether $\Delta^{\text{val}}_{Y_\bullet}(D)$ is a numerical invariant of D or not. If D is a big divisor on X, then by Remark 3.9, $\Delta^{\text{val}}_{Y_\bullet}(D)$ depends only on the numerical class of D. However, in general, we may have $\kappa(D) < \kappa(D')$ for divisors D, D' satisfying $D \equiv D'$ (see [Le, Example 6.1]). Thus by choosing an admissible flag $Y_\bullet$ containing a Nakayama subvariety of both D and D', we obtain $\Delta^{\text{val}}_{Y_\bullet}(D) \neq \Delta^{\text{val}}_{Y_\bullet}(D')$.

Remark 3.14. If we drop the condition $\kappa(D) \geq 0$, then Theorem 3.12 does not hold. For example, take two distint points P, Q on $\mathbb{P}^1$ and an irrational number a as in [G, Remark 2.1]. Set $D := a(P - Q)$. Then $D \sim_{\mathbb{R}} 0$ and $\Delta^{\text{val}}_{Y_\bullet}(D) = \{0 \in \mathbb{R}^1\}$. However, we have $\kappa(D) = -\infty$.

On the other hand, we have the following.

Proposition 3.15. *If two effective divisors D, D' satisfy $D \sim_{\mathbb{R}} D'$, then $\Delta^{\text{val}}_{Y_\bullet}(D) = \Delta^{\text{val}}_{Y_\bullet}(D')$ with respect to any admissible flag $Y_\bullet$.*

Proof. It is trivial. □

Remark 3.16. The converse of Proposition 3.15 is false in general. For example, consider a minimal ruled surface $f : S \to C$ over a smooth projective curve of genus $g \geq 1$. Let $F_1 := f^*P$ and $F_2 := f^*Q$ where P and Q are two distinct points on C. Note that $F_1 \not\sim_{\mathbb{R}} F_2$. However, we claim that $\Delta^{\text{val}}_{Y_\bullet}(F_1) = \Delta^{\text{val}}_{Y_\bullet}(F_2)$ for any admissible flag $Y_\bullet$ on S. To see this, observe that a curve Y_1 is either a fiber of f or dominating C via f. In the first case, we have

$$\Delta^{\text{val}}_{Y_\bullet}(F_1) = \Delta^{\text{val}}_{Y_\bullet}(F_2) = \{(x_1, x_2) \in \mathbb{R}^2 \mid 0 \leq x_1 \leq 1, x_2 = 0\}.$$

In the second case, note that Y_1 is a Nakayama subvariety of both F_1 and F_2. Furthermore, we have

$$\mathrm{vol}_{S|Y_1}(F_1) = Y_1 \cdot F_1 = Y_1 \cdot F_2 = \mathrm{vol}_{S|Y_1}(F_2),$$

and hence, the claim follows.

3.3. Limiting Okounkov body $\Delta^{\lim}_{Y_\bullet}(D)$.

We now define and study the limiting Okounkov body $\Delta^{\lim}_{Y_\bullet}(D)$ of a pseudoeffective divisor D on a variety X of dimension n.

Definition 3.17. Let D be a pseudoeffective divisor D on a variety X of dimension n. The *limiting Okounkov body* $\Delta^{\lim}_{Y_\bullet}(D)$ of D with respect to an admissible flag $Y_\bullet$ is defined as

$$\Delta^{\lim}_{Y_\bullet}(D) := \lim_{\varepsilon \to 0+} \Delta_{Y_\bullet}(D + \varepsilon A) \subseteq \mathbb{R}^n$$

where A is any ample divisor. If D is not a pseudoeffective divisor, then we define $\Delta^{\lim}(D) := \emptyset$.

Remark 3.18. It is easy to see that the definition of the limiting Okounkov body does not depend on the choice of an ample divisor A. If D is big, then $\Delta^{\lim}_{Y_\bullet}(D)$ coincides with $\Delta_{Y_\bullet}(D)$ for any admissible flag $Y_\bullet$ on X.

We are mainly interested in the limiting Okounkov bodies $\Delta^{\lim}_{Y_\bullet}(D)$ when $Y_\bullet$ contains contains a positive volume subvariety V of D. To investigate such $\Delta^{\lim}_{Y_\bullet}(D)$ in detail, we introduce the following alternative construction, which also gives the same convex body as $\Delta^{\lim}_{Y_\bullet}(D)$ after all (see Proposition 3.21). This construction is often more convenient to study the asymptotic invariants of pseudoeffective divisors. To begin with, we first consider a pseudoeffective Cartier divisor D on a variety X of dimension n. Let $V \subseteq X$ be a positive volume subvariety of D. Fix an admissible flag $V_\bullet$ on V:

$$V_\bullet : V = V_0 \supseteq V_1 \supseteq \cdots \supseteq V_{\kappa_\nu(D)} = \{x\}.$$

Let A be an ample Cartier divisor on X. For each integer $k \geq 1$, consider the restricted graded linear series $W^k_\bullet := W_\bullet(kD + A|V)$ of $kD + A$ along V given by $W_m(kD + A|V) = h^0(X|V, m(kD + A))$ for $m \geq 0$. We define the *restricted limiting Okounkov body* of a Cartier divisor D with respect to a positive volume subvariety V of D as

$$\Delta^{\lim}_{V_\bullet}(D) := \lim_{k \to \infty} \frac{1}{k} \Delta_{V_\bullet}(W^k_\bullet) \subseteq \mathbb{R}^{\kappa_\nu(D)}.$$

By the continuity, we can extend this definition for any pseudoeffective $\mathbb{R}$-divisor D.

Definition 3.19. Let D be a pseudoeffective divisor D on a variety X with a positive volume subvariety $V \subseteq X$. The *restricted limiting Okounkov body* $\Delta^{\lim}_{V_\bullet}(D)$ of D with respect to $V_\bullet$ as

$$\Delta^{\lim}_{V_\bullet}(D) := \lim_{\varepsilon \to 0+} \Delta_{V_\bullet}(D + \varepsilon A) \subseteq \mathbb{R}^{\kappa_\nu(D)}$$

for some ample divisor A on X. If D is not pseudoeffective, then we define $\Delta^{\lim}_{V_\bullet}(D) := \emptyset$.

This definition depends on the choice of V and the admissible flag $V_\bullet$ on it, but it is independent of the choice of A. By definition, $\Delta^{\lim}_{V_\bullet}(D)$ is a closed convex subset of $\mathbb{R}^{\kappa_\nu(D)}$. By the inclusion $\mathbb{R}^{\kappa_\nu(D)} = \{0\}^{n-\kappa_\nu(D)} \times \mathbb{R}^{\kappa_\nu(D)} \hookrightarrow \mathbb{R}^n$, we often treat $\Delta^{\lim}_{V_\bullet}(D)$ as a subset of $\mathbb{R}^n$.

The following is the main property of the limiting Okounkov body $\Delta^{\lim}_{V_\bullet}(D)$.

Theorem 3.20. *Let D be a pseudoeffective divisor on a variety X. Fix a positive volume subvariety $V \subseteq X$ of D and consider an admissible flag $V_\bullet$ on V. Then we have*

$$\dim \Delta^{\lim}_{V_\bullet}(D) = \kappa_\nu(D) \quad \textit{and} \quad \mathrm{vol}_{\mathbb{R}^{\kappa_\nu(D)}}(\Delta^{\lim}_{V_\bullet}(D)) = \frac{1}{\kappa_\nu(D)!} \mathrm{vol}^+_{X|V}(D).$$

Proof. It is enough to consider the case where D is a Cartier divisor since the $\mathbb{R}$-divisor case follows by the standard argument. Since $V \not\subseteq \mathbf{B}_-(D)$ by definition, it follows that $V \not\subseteq \mathbf{B}_+(kD+A)$ for any $k \geq 1$. Let $W^k_\bullet := W_\bullet(kD + A|V)$ be the restricted graded series of $kD + A$ along V for $k \geq 1$. Then $W^k_\bullet$ satisfies Condition (C) by [LM, Lemma 2.16]. It follows from Theorem 3.5 that for each $k \geq 1$, we have

$$\dim \Delta_{V_\bullet}(W^k_\bullet) = \dim V = \kappa_\nu(D)$$

and

$$\mathrm{vol}_{\mathbb{R}^{\kappa_\nu(D)}}(\Delta_{V_\bullet}(W^k_\bullet)) = \frac{1}{\kappa_\nu(D)!}\,\mathrm{vol}_{X|V}(kD + A).$$

The first equality implies $\dim \Delta^{\lim}_{V_\bullet}(D) = \kappa_\nu(D)$, and the second equality yields the following computation:

$$\begin{aligned}
\mathrm{vol}_{\mathbb{R}^{\kappa_\nu(D)}}(\Delta^{\lim}_{V_\bullet}(D)) &= \mathrm{vol}_{\mathbb{R}^{\kappa_\nu(D)}}\left(\lim_{k\to\infty} \tfrac{1}{k}\Delta_{V_\bullet}(W^k_\bullet)\right) \\
&= \lim_{k\to\infty} \tfrac{1}{k^{\kappa_\nu(D)}}\,\mathrm{vol}_{\mathbb{R}^{\kappa_\nu(D)}}\,\Delta_{V_\bullet}(W^k_\bullet) \\
&= \lim_{k\to\infty} \tfrac{1}{\kappa_\nu(D)!}\,\mathrm{vol}_{X|V}\left(D + \tfrac{1}{k}A\right) \\
&= \tfrac{1}{\kappa_\nu(D)!}\,\mathrm{vol}^+_{X|V}(D).
\end{aligned}$$

□

Proposition 3.21. *Let D be a pseudoeffective divisor on a variety X of dimension n, and fix an admissible flag $Y_\bullet$ containing a positive volume subvariety $V = Y_{n-\kappa_\nu(D)}$ of D. Then*

$$\Delta^{\lim}_{Y_\bullet}(D) = \Delta^{\lim}_{V_\bullet}(D).$$

In particular, $\Delta^{\lim}_{Y_\bullet}(D)$ depends on the $(n - \kappa_\nu(D))$-th partial flag $V_\bullet$.

Proof. We will denote $\kappa_\nu := \kappa_\nu(D)$. If $\kappa_\nu = n$, then there is nothing to prove. Therefore, we assume that $0 \leq \kappa_\nu < n$. Fix an ample divisor A on X. By definition, the positive volume subvariety $Y_{n-\kappa_\nu} = V$ is not contained in $\mathbf{B}_-(D)$. Since $Y_i \supseteq Y_{n-\kappa_\nu}$ for $0 \leq i \leq n - \kappa_\nu$, we have $Y_i \not\subseteq \mathbf{B}_+(D + \varepsilon A)$ for any $\varepsilon > 0$. Thus [LM, Theorem 4.26] implies that if $0 \leq i \leq n - \kappa_\nu$, then $\Delta_{Y_\bullet}(D + \varepsilon A) \cap (\{0\}^i \times \mathbb{R}^{n-i}) = \Delta_{Y_{i\bullet}}(D + \varepsilon A)$. By taking the limit $\varepsilon \to 0$, we obtain for $0 \leq i \leq n - \kappa_\nu$

$$(*) \qquad \Delta^{\lim}_{Y_\bullet}(D) \cap (\{0\}^i \times \mathbb{R}^{n-i}) = \lim_{\varepsilon\to 0+} \Delta_{Y_{i\bullet}}(D + \varepsilon A) =: \Delta^{\lim}_{Y_{i\bullet}}(D).$$

Since $Y_{n-\kappa_\nu} = V$, it is enough to check $\Delta^{\lim}_{Y_\bullet}(D) \subseteq \{0\}^{n-\kappa_\nu} \times \mathbb{R}^{\kappa_\nu}$ to prove $\Delta^{\lim}_{Y_\bullet}(D) = \Delta^{\lim}_{V_\bullet}(D)$. Suppose that this does not hold, i.e.,

$$\Delta^{\lim}_{Y_\bullet}(D) \not\subseteq \{0\}^{n-\kappa_\nu} \times \mathbb{R}^{\kappa_\nu} \subseteq \mathbb{R}^n.$$

Then by Theorem 3.20, $l := \dim \Delta^{\lim}_{Y_\bullet}(D) > \kappa_\nu = \dim \Delta^{\lim}_{V_\bullet}(D)$ and we have

$$\Delta^{\lim}_{Y_\bullet}(D) \subseteq \mathbb{R}^{i_1} \times \mathbb{R}^{i_2} \times \cdots \times \mathbb{R}^{i_{n-\kappa_\nu}} \times \mathbb{R}^{\kappa_\nu}$$

where $i_j = 0$ or 1, $\sum_{j=1}^{n-\kappa_\nu} i_j = l - \kappa_\nu$ and $\mathbb{R}^0 := \{0\}$. For simplicity, we may just assume

$$\Delta^{\lim}_{Y_\bullet}(D) \subseteq \mathbb{R}^{l-\kappa_\nu} \times \{0\}^{n-l} \times \mathbb{R}^{\kappa_\nu}.$$

Now consider a sufficiently positive divisor A' so that $\Delta_{Y_\bullet}(\varepsilon A')$ contains n-simplex $\triangle_\varepsilon$ of length ε:

$$\triangle_\varepsilon := \left\{ (x_1, \cdots, x_n) \,\middle|\, x_i \geq 0 \text{ and } \sum_{i=1}^n x_i \leq \varepsilon \right\} \subseteq \Delta_{Y_\bullet}(\varepsilon A')$$

for all $\varepsilon > 0$. We may also assume that A is sufficiently ample so that $A'' := A - A'$ is also ample. By the convexity of $\Delta_{Y_\bullet}(D)$ (cf. [LM, Proof of Corollary 4.12]), we have

$$\Delta^{\lim}_{Y_\bullet}(D) + \triangle_\varepsilon \subseteq \Delta^{\lim}_{Y_\bullet}(D) + \Delta_{Y_\bullet}(\varepsilon A') \subseteq \Delta_{Y_\bullet}(D + \varepsilon A'') + \Delta_{Y_\bullet}(\varepsilon A') \subseteq \Delta_{Y_\bullet}(D + \varepsilon A).$$

Thus by considering the Euclidean volumes in $\mathbb{R}^n$ of the convex hull of $\Delta^{\lim}_{Y_\bullet}(D) \cup \triangle_\varepsilon \subseteq \Delta^{\lim}_{Y_\bullet}(D) + \triangle_\varepsilon$ and $\Delta_{Y_\bullet}(D+\varepsilon A)$, we obtain

$$C \cdot \mathrm{vol}_{\mathbb{R}^l} \Delta^{\lim}_{Y_\bullet}(D) \cdot \varepsilon^{n-l} \leq \mathrm{vol}_{\mathbb{R}^n} \Delta_{Y_\bullet}(D+\varepsilon A) = \frac{1}{n!} \cdot \mathrm{vol}(D+\varepsilon A).$$

for some constant $C > 0$ depending only on n and l. Consequently, $C' \cdot \varepsilon^{n-l} \leq \mathrm{vol}(D+\varepsilon A)$ for the constant $C' := C \cdot n! \cdot \mathrm{vol}_{\mathbb{R}^l} \Delta^{\lim}_{Y_\bullet}(D) > 0$. However, this is a contradiction to [Le, Theorem 1.1 (2)] since $l > \kappa_\nu$. ☐

Corollary 3.22. *Let D be a pseudoeffective divisor on a smooth projective variety X. Fix a positive volume subvariety $V \subseteq X$ of D and consider an admissible flag $Y_\bullet$ on X containing V. Then we have*

$$\dim \Delta^{\lim}_{Y_\bullet}(D) = \kappa_\nu(D) \quad \textit{and} \quad \mathrm{vol}_{\mathbb{R}^{\kappa_\nu(D)}}(\Delta^{\lim}_{Y_\bullet}(D)) = \frac{1}{\kappa_\nu(D)!} \mathrm{vol}^+_{X|V}(D).$$

Proof. Immediate by Theorem 3.20 and Proposition 3.21. ☐

Remark 3.23. We do not need to assume the generality of $Y_n = \{x\}$ as we did for the valuative Okounkov body in Theorem 3.12.

Remark 3.24. Boucksom also studies the limiting Okounkov body (the numerical Okounkov body in his terminology) of a pseudoeffective divisor with respect to any admissible flag in [B, 4.1.3]. He defines the limiting Okounkov body as a fiber of the global Okounkov body (see [LM, Theorem B]) over a given pseudoeffective divisor class. He proves that $\dim \Delta^{\lim}_{Y_\bullet}(D) \leq \kappa_\nu(D)$ for any admissible flag $Y_\bullet$ ([B, Lemma 4.8]) and the inequality can be strict ([B, Example 4.14]). Furthermore, he also shows that there exists an admissible flag $Y_\bullet$ such that $\dim \Delta^{\lim}_{Y_\bullet}(D) = \kappa_\nu(D)$ when D is nef ([B, Proposition 4.9]). Our result Corollary 3.22 can be regarded as a generalization of his result.

Using Corollary 3.22, we can give a new characterization of numerical Iitaka dimension.

Corollary 3.25. *Let D be a pseudoeffective divisor on a smooth projective variety X. Then we have*

$$\kappa_\nu(D) = \max\{\dim \Delta^{\lim}_{Y_\bullet}(D) \mid Y_\bullet \textit{ is an admissible flag on } X\ \}.$$

Proof. It follows from Corollary 3.22 and [B, Lemma 4.8]. ☐

If D is a big divisor, then by [LM, Proposition 4.1] and [J, Theorem A] it is known that $\Delta_{Y_\bullet}(D) = \Delta_{Y_\bullet}(D')$ for all admissible flags $Y_\bullet$ on X if and only if $D \equiv D'$. Remark 3.13 implies that the statement is false in general in the pseudoeffective case for $\Delta_{Y_\bullet}(D)$ (as well as $\Delta^{\mathrm{val}}_{Y_\bullet}(D)$). We prove next that $\Delta^{\lim}_{Y_\bullet}(D)$ is an appropriate generalization of $\Delta_{Y_\bullet}(D)$ which makes the statement true. We first consider a special case.

Proposition 3.26. *Let D be a pseudoeffective divisor on a smooth projective variety X. Fix a positive volume subvariety $V \subseteq X$ of D and consider an admissible flag $Y_\bullet$ on X containing V. Then $\Delta^{\lim}_{Y_\bullet}(D)$ depends on the numerical class of D.*

Proof. By Proposition 3.21, we only have to prove the assertion for the $V_\bullet$ instead of $Y_\bullet$. Suppose that D, D' are pseudoeffective divisors such that $D \equiv D'$. Then Theorem 2.18 implies that $(D+\varepsilon A)|_V$, $(D'+\varepsilon A)|_V$ are big on V for a fixed ample divisor A and any $\varepsilon > 0$. By Remark 3.7, we have $\Delta_{V_\bullet}(D+\varepsilon A) = \Delta_{V_\bullet}(D'+\varepsilon A)$ for any admissible flags $V_\bullet$ on V. Thus we get

$$\Delta^{\lim}_{V_\bullet}(D) = \lim_{\varepsilon \to 0+} \Delta_{V_\bullet}(D+\varepsilon A) = \lim_{\varepsilon \to 0+} \Delta_{V_\bullet}(D'+\varepsilon A) = \Delta^{\lim}_{V_\bullet}(D').$$

☐

For a pseudoeffective divisor D, we define the following set of divisors

$$div(\Delta^{\lim}(D)) := \left\{ E \,\middle|\, \begin{array}{l} \text{there exists an admissible flag } Y_\bullet \text{ with } Y_1 = E \\ \text{such that} \Delta^{\lim}_{Y_\bullet}(D)_{x_1=0} = \emptyset \end{array} \right\}.$$

Lemma 3.27. *The divisors in $div(\Delta^{\lim}(D))$ are precisely the divisorial components of $\mathbf{B}_-(D)$ and the set $div(\Delta^{\lim}(D))$ is finite.*

Proof. Let $Y_\bullet$ be an admissible flag on X such that $Y_1 = E$. By the definition of the Okounkov body, it is easy to see that for any ample divisor A and $\varepsilon > 0$,

$$\operatorname{ord}_E(||D + \varepsilon A||) = \inf \{x_1 \,|(x_1, \cdots, x_n) \in \Delta_{Y_\bullet}(D + \varepsilon A)\}.$$

If we let $\varepsilon \to 0$, then we have

$$\operatorname{ord}_E(||D||) = \inf \left\{ x_1 \,\middle|(x_1, \cdots, x_n) \in \Delta^{\lim}_{Y_\bullet}(D) \right\}.$$

By Theorem 2.1, $\operatorname{ord}_E(||D||) > 0$ if and only if $E \subseteq \mathbf{B}_-(D)$. Thus we obtain the desired statement. The set $div(\Delta^{\lim}(D))$ is finite by [N, Corollary 1.11]. □

We now prove a generalization of Jow's theorem for the limiting Okounkov bodies.

Theorem 3.28. *Let D and D' be pseudoeffective divisors on X. Then the following are equivalent:*

(1) $D \equiv D'$

(2) $\Delta^{\lim}_{Y_\bullet}(D) = \Delta^{\lim}_{Y_\bullet}(D')$ *with respect to any admissible flags $Y_\bullet$ on X.*

Proof. The equivalence (1) ⇔ (2) is known for big divisors by [LM, Proposition 4.1] and [J, Theorem A]. Thus we assume that D, D' are not big. The implication (1)⇒(2) follows from the big case: since $\Delta_{Y_\bullet}(D + \varepsilon A) = \Delta_{Y_\bullet}(D' + \varepsilon A)$ for any ample divisor A and any $\varepsilon > 0$, we have

$$\Delta^{\lim}_{Y_\bullet}(D) = \lim_{\varepsilon \to 0} \Delta_{Y_\bullet}(D + \varepsilon A) = \lim_{\varepsilon \to 0} \Delta_{Y_\bullet}(D' + \varepsilon A) = \Delta^{\lim}_{Y_\bullet}(D').$$

To prove (1)⇐(2), we use the argument used to prove the big case by Jow in [J]. Let $E_1, \cdots, E_l$ be the divisorial components of $\mathbf{B}_-(D)$ and A an ample divisor on X. For any sufficiently general admissible flag $Y_\bullet$ on X such that $Y_{n-1} \not\subseteq \mathbf{B}_-(D)$, we have $Y_{n-1} \not\subseteq \mathbf{B}_+(D + A)$ for any ample divisor A and

$$\begin{aligned} & \operatorname{vol}_{\mathbb{R}}(\Delta^{\lim}_{Y_\bullet}(D)_{x_1=\cdots=x_{n-1}=0}) \\ = \; & \operatorname{vol}_{\mathbb{R}} \left(\lim_{\varepsilon \to 0+} \Delta_{Y_\bullet}(D + \varepsilon A) \right)_{x_1=\cdots=x_{n-1}=0} \\ = \; & \lim_{\varepsilon \to 0+} \operatorname{vol}_{\mathbb{R}}(\Delta_{Y_\bullet}(D + \varepsilon A)_{x_1=\cdots=x_{n-1}=0}) \\ = \; & \lim_{\varepsilon \to 0+} \operatorname{vol}_{X|Y_{n-1}}(D + \varepsilon A) \qquad \text{(by [J, Theorem 3.4 (b)])} \\ = \; & Y_{n-1} \cdot D - \sum_{i=1}^{l} \sum_{p \in Y_{n-1} \cap E_i} \operatorname{ord}_{E_i}(||D||) \qquad \text{(by Lemma 3.29).} \end{aligned}$$

As can be seen in the proof of Lemma 3.27, we can read off $\operatorname{ord}_{E_i}(||D||)$ from the limiting Okounkov bodies $\Delta^{\lim}_{Y'_\bullet}(D)$ with respect to admissible flags $Y'_\bullet$ such that $Y'_1 = E_i$. Therefore, if $\Delta^{\lim}_{Y_\bullet}(D) = \Delta^{\lim}_{Y_\bullet}(D')$ for any admissible flags $Y_\bullet$ on X, then we have $Y_{n-1} \cdot D = Y_{n-1} \cdot D'$. As in [J], if we now consider sufficiently general flags $Y^1_\bullet, \cdots, Y^\rho_\bullet$ (where $\rho = \dim \mathrm{N}^1(X)_{\mathbb{R}}$) such that $Y^1_{n-1}, \cdots, Y^\rho_{n-1}$ form a basis of $\mathrm{N}_1(X)_{\mathbb{R}}$, then we can conclude that $D \equiv D'$. □

It remains to prove the following lemma which is used in the proof above.

Lemma 3.29 (cf. [J, Corollary 3.3])**.** *Let $Y_\bullet$ be a sufficiently general admissible flag on X. For a pseudoeffective divisor D on X, let $E_1, E_2, \cdots, E_l$ be the divisorial components of $\mathbf{B}_-(D)$. Then we have*

$$\operatorname{vol}^+_{X|Y_{n-1}}(D) = \lim_{\varepsilon\to 0+} \operatorname{vol}_{X|Y_{n-1}}(D+\varepsilon A) = Y_{n-1}\cdot D - \sum_{i=1}^{l} \sum_{p\in Y_{n-1}\cap E_i} \operatorname{ord}_{E_i}(||D||).$$

Proof. Suppose first that D is big. Since Y_{n-1} is very general, we have $D \in \operatorname{Big}^{Y_{n-1}}(X)$. Thus $\operatorname{vol}^+_{X|Y_{n-1}}(D) = \operatorname{vol}_{X|Y_{n-1}}(D)$ and the statement is nothing but [J, Corollary 3.3].

Now let D be a pseudoeffective divisor. Applying the statement for the case of big divisors to $\operatorname{vol}_{X|Y_{n-1}}(D+\varepsilon A)$, we obtain

$$\begin{aligned}\operatorname{vol}^+_{X|Y_{n-1}}(D) &= \lim_{\varepsilon\to 0+} \operatorname{vol}_{X|Y_{n-1}}(D+\varepsilon A)\\ &= \lim_{\varepsilon\to 0+}\left(Y_{n-1}\cdot (D+\varepsilon A) - \sum_{i=1}^{l}\sum_{p\in Y_{n-1}\cap E_i} \operatorname{ord}_{E_i}(||D+\varepsilon A||)\right)\\ &= Y_{n-1}\cdot D - \sum_{i=1}^{l}\sum_{p\in Y_{n-1}\cap E_i} \operatorname{ord}_{E_i}(||D||).\end{aligned}$$

□

Remark 3.30. Fulger pointed out to us that in the equation of Lemma 3.29 we actually have

$$\sum_{i=1}^{l}\sum_{p\in Y_{n-1}\cap E_i} \operatorname{ord}_{E_i}(||D||) = Y_{n-1}\cdot N$$

where N is the negative part of the divisorial Zariski decomposition of D.

Remark 3.31. In Theorem 3.28, it is tempting to expect that $D \equiv D'$ if and only if $\Delta^{\lim}_{Y_\bullet}(D) = \Delta^{\lim}_{Y_\bullet}(D')$ for any admissible flags $Y_\bullet$ containing positive volume subvarieties of both D and D'. However, it is not true and we also need to consider the admissible flags not containing the positive volume loci of the divisor. For example, consider a pseudoeffective divisor $D \not\equiv 0$ on a variety X with $\kappa_\nu(D) = 0$. Let $D' := 2D$ so that $D \not\equiv D'$. Every point $x \in X$ with $x \notin \mathbf{B}_-(D) = \mathbf{B}_-(D')$ is a positive volume subvariety of both D and D'. However, we see that for any admissible flag $Y_\bullet$ containing x, we have

$$\Delta^{\lim}_{Y_\bullet}(D) = \Delta^{\lim}_{Y_\bullet}(D') = \{\text{the origin of } \mathbb{R}^{\dim X}\}.$$

In fact, if we consider the admissible flags $Y_\bullet$ containing positive volume subvarieties of both D and D' in Theorem 3.28, then using Remark 3.30 we only obtain $P \equiv P'$ where $D = P + N$ and $D' = P' + N'$ are the divisorial Zariski decompositions.

Remark 3.32. Let D be a pseudoeffective Cartier divisor on X and $V \subseteq X$ be its positive volume subvariety of dimension $\kappa_\nu(D)$. Let $Y_\bullet$ be an admissible flag on X whose $(n-\kappa_\nu(D))$-th partial flag $Y_{(n-\kappa_\nu(D))\bullet}$ induces an admissible flag $V_\bullet$ on V. To construct the limiting Okounkov body $\Delta^{\lim}_{V_\bullet}(D)$, we used the the restricted complete linear series $W_\bullet(X|V, D)$ which is a subseries of the complete linear series $W'_\bullet = \{H^0(V, \mathcal{O}_V(mD|_V))\}_{m\geq 0}$. Using the linear series $W'_\bullet$ instead, we can construct a convex body $\Delta_{Y_{V_\bullet}}(D|_V)$ associated to the restricted divisor $D|_V$ on V. We can easily verify the following:

$$\dim \Delta_{V_\bullet}(D|_V) = \kappa_\nu(D) \text{ and } \operatorname{vol}_{\mathbb{R}^{\kappa_\nu(D)}} \Delta_{V_\bullet}(D|_V) = \frac{1}{\kappa_\nu(D)!}\operatorname{vol}_V(D|_V).$$

The proofs are left to the readers. Let $Y_\bullet$ be an admissible flag on X which is general enough so that it contains both a Nakayama subvariety U and a positive volume subvariety V of a pseudoeffective divisor D. Then we have the following inclusions:

$$(\#) \qquad \Delta^{\text{val}}_{Y_\bullet}(D) \subseteq \Delta^{\lim}_{Y_\bullet}(D) = \Delta^{\lim}_{V_\bullet}(D) \subseteq \Delta_{V_\bullet}(D|_V).$$

This confirms the inclusions $\text{vol}_{X|V}(D) \leq \text{vol}^+_{X|V}(D) \leq \text{vol}_V(D|_V)$ which we saw in Subsection 2.3. If D is big, then all the inclusions are equalities. However, we will see in the next section that if D is not big, then they are strict in general.

4. EXAMPLES

In this section, we exhibit various examples and counterexamples related to our results. The following example shows that for some badly chosen flags, Theorems 3.12 and 3.20 do not hold.

Example 4.1. Let $\pi : S \to \mathbb{P}^2$ be a blow-up of $\mathbb{P}^2$ at two distinct points with exceptional divisors E_1 and E_2 and $L := \pi^*\mathcal{O}_{\mathbb{P}^2}(1)$. Consider a non-big effective divisor $D := L - E_1 + E_2$, and fix an admissible flag

$$Y_\bullet : S \supseteq E_1 \supseteq \{x\}.$$

Here we note that E_1 is neither a Nakayama subvariety nor a positive volume subvariety of D. We have

$$\Delta^{\text{val}}_{Y_\bullet}(D) = \Delta^{\lim}_{Y_\bullet}(D) = \{(x_1, x_2) \in \mathbb{R}^2 \mid 1 \leq x_1 \leq 2 \text{ and } x_2 = 0\}$$

whose Euclidean volume in the x_1-axis is 1. However, note that $\mathbf{B}_-(D) = \mathbf{B}_+(D + \varepsilon A) = E_2$ for an ample divisor A and a small $\varepsilon > 0$. Thus we see that $\text{vol}_{X|E_1}(D) = \text{vol}^+_{X|E_1}(D) = 0$. We also remark that $\Delta^{\text{val}}_{Y_\bullet}(D) = \Delta^{\lim}_{Y_\bullet}(D)$ is not in the x_2-axis as usual.

Examples 4.2 and 4.3 show that the inequalities in (#) of Remark 3.32 can be strict.

Example 4.2. In this example, we see that $\Delta^{\text{val}}_{Y_\bullet}(D) \subsetneq \Delta^{\lim}_{Y_\bullet}(D)$ even if $\kappa(D) = \kappa_\nu(D)$. Let S be a relatively minimal rational surface, and H a sufficiently positive ample divisor. Take a general element $V \in |H|$. Fix an admissible flag

$$Y_\bullet : S \supseteq V \supseteq \{x\}.$$

Note that $\kappa(-K_S) = \kappa_\nu(-K_S) = 1$, and V is both a Nakayama subvariety and a positive volume subvariety of D. Thus we have $\text{vol}_{\mathbb{R}^1}(\Delta^{\text{val}}_{Y_\bullet}(-K_S)) = \text{vol}_{S|V}(-K_S)$ and $\text{vol}_{\mathbb{R}^1}(\Delta^{\lim}_{Y_\bullet}(-K_S)) = \text{vol}^+_{S|V}(-K_S)$. However, we saw in Example 2.3 that $\text{vol}_{S|V}(-K_S) < \text{vol}^+_{S|V}(-K_S)$. Thus $\Delta^{\text{val}}_{Y_\bullet}(-K_S) \subsetneq \Delta^{\lim}_{Y_\bullet}(-K_S)$.

Example 4.3. Let $S := \mathbb{P}(E)$ where E is a rank two vector bundle on an elliptic curve C such that it is a nontrivial extension of $\mathcal{O}_C$ by $\mathcal{O}_C$, and H be the tautological divisor of $\mathbb{P}(E)$. Then we can easily check that H is nef and $\kappa(H) = 0$ but $\kappa_\nu(H) = 1$. Let F be a fiber of the natural ruling $\pi : S \to C$. Note that any point in S is a Nakayama subvariety of H and F is a positive volume subvariety of H. Thus take an admissible flag

$$Y_\bullet : S \supseteq F \supseteq \{x\}.$$

Then we can easily compute the following.

(1) $\Delta^{\text{val}}_{Y_\bullet}(H) = \{(0,0)\}$ and $\text{vol}_{S|\{x\}}(H) = 1$.
(2) $\Delta^{\lim}_{Y_\bullet}(H) = \{(x_1, x_2) \mid x_1 = 0 \text{ and } 0 \leq x_2 \leq 1\}$ and $\text{vol}^+_{S|F}(H) = 1$.
(3) $\Delta_{Y_\bullet}(H|_F) = \Delta^{\lim}_{Y_\bullet}(H)$.

For $\Delta^{\lim}_{Y_\bullet}(H)$, we see the convergence of $\lim_{\varepsilon \to 0+} \Delta_{Y_\bullet}(H + \varepsilon A)$ for any ample divisor A in the following picture.



In fact, we use the following theorem to compute the limiting Okounkov body $\Delta^{\lim}_{Y_\bullet}(D)$ of a pseudoeffective divisor D on a surface (cf. [LM, Theorem 6.4])

Theorem 4.4. *Let S be a smooth projective surface, and D be a pseudoeffective divisor on S. Fix an admissible flag $Y_\bullet : S \supseteq C \supseteq \{x\}$. Let $a := \operatorname{mult}_C N$ where $D = P + N$ is the Zariski decomposition, and $\mu := \sup\{s \geq 0 \mid D - sC$ is pseudoeffective$\}$. Consider a divisor $D_t := D - tC$ for $a \leq t \leq \mu$. Denote by $D_t = P_t + N_t$ the Zariski decomposition. Let $\alpha(t) := \operatorname{ord}_x(N_t|_C)$ and $\beta(t) := \alpha(t) + C.P_t$. Then the limiting Okounkov body of D is given by*

$$\Delta^{\lim}_{Y_\bullet}(D) = \{(x_1, x_2) \in \mathbb{R}^2 \mid a \leq x_1 \leq \mu \text{ and } \alpha(x_1) \leq x_2 \leq \beta(x_2)\}.$$

Proof. Let $D^\varepsilon := D + \varepsilon A$ for some ample divisor A, and $D^\varepsilon_t := D^\varepsilon - tC$. Denote by $D^\varepsilon_t = P^\varepsilon_t + N^\varepsilon_t$ the Zariski decomposition. We see that the Zariski decomposition $D_t = P_t + N_t$ is given by $P_t = \lim_{\varepsilon \to 0+} P^\varepsilon_t$ and $N_t = \lim_{\varepsilon \to 0+} N^\varepsilon_t$. Since $\Delta^{\lim}_{Y_\bullet}(D) = \lim_{\varepsilon \to 0+} \Delta_{Y_\bullet}(D + \varepsilon A)$, the assertion now follows from [LM, Theorem 6.4]. □

Remark 4.5. In Theorem 4.4, we do not need to assume that an admissible flag $Y_\bullet$ contains a positive volume subvariety.

Lastly, we give an example for which the inequalities (#) of Remark 3.32 are equalities.

Example 4.6. Let X be a smooth projective toric variety with a torus T. For definitions and notations for toric varieties, we refer to [F]. When D is a big T-divisor, $\Delta^{\lim}_{Y_\bullet}(D) = \Delta^{\mathrm{val}}_{Y_\bullet}(D) = \Delta_{Y_\bullet}(D)$ is the same as P_D up to translation for any fixed admissible flag $Y_\bullet$. Now let D be a pseudoeffective T-divisor on X. Then we have

$$\Delta^{\lim}_{Y_\bullet}(D) = \lim_{\varepsilon \to 0+} \Delta_{Y_\bullet}(D + \varepsilon A) = \bigcap_{\varepsilon > 0} \Delta_{Y_\bullet}(D + \varepsilon A) = \bigcap_{\varepsilon > 0} P_{D + \varepsilon A} = P_D.$$

For any smooth projective toric variety, we have $\kappa(D) = \kappa_\nu(D)$ so that $\dim \Delta^{\mathrm{val}}_{Y_\bullet}(D) = \dim \Delta^{\lim}_{Y_\bullet}(D)$. In fact, we can show that $\Delta^{\mathrm{val}}_{Y_\bullet}(D) = \Delta^{\lim}_{Y_\bullet}(D)$ for some admissible flag $Y_\bullet$. We first observe that there is a torus invariant subvariety of X which is both a Nakayama subvariety of D and a positive volume subvariety of D. To see this, consider a rational map $f : X \dashrightarrow Y$ given by the linear system $|\lfloor mD \rfloor|$ for $m \gg 0$. Then any torus invariant subvariety V of X dominating Y via f is such a special subvariety. Now assume that an admissible flag $Y_\bullet$ contains V. We now show that $\Delta^{\mathrm{val}}_{Y_\bullet}(D) = \Delta^{\lim}_{Y_\bullet}(D)$. It is enough to show that $\operatorname{vol}_{X|V}(D) = \operatorname{vol}^+_{X|V}(D) = \lim_{\varepsilon \to 0+} \operatorname{vol}_{X|V}(D + \varepsilon A)$. Let $V(\tau)$ be a T-invariant irreducible closed subvariety of X, and π the projection of $M_{\mathbb{R}}$ to $(M_\tau)_{\mathbb{R}} = M_{\mathbb{R}} \cap \tau^\perp$. Then we observe that

(1) $\dim H^0(X, \mathcal{O}_X(D)) =$ the number of lattice points in P_D, and
(2) $\dim \operatorname{Im}(H^0(X, \mathcal{O}_X(D)) \to H^0(V, \mathcal{O}_V(D|_V))) =$ the number of lattice points in $\pi(P_D)$.

It follows that $\pi(P_D) = \bigcap_{\varepsilon > 0} \pi\left(P_{D + \varepsilon A}\right)$. Thus $\operatorname{vol}_{X|V}(D) = \operatorname{vol}^+_{X|V}(D)$.

References

[B] S. Boucksom, *Corps D'Okounkov (d'après Okounkov, Lazarsfeld- Mustaţă et Kaveh-Khovanskii)*, Séminaire Bourbaki, 65ème année, 2012-2013, no. 1059.

[BBP] S. Boucksom, A. Broustet, and G. Pacienza, *Uniruledness of stable base loci of adjoint linear systems via Mori Theory*, Math. Z. **275** (2013), 499-507.

[ELMNP1] L. Ein, R. Lazarsfeld, M. Mustaţă, M. Nakamaye, and M. Popa, *Asymptotic invariants of base loci*, Ann. Inst. Fourier, **56** (2006), 1701-1734.

[ELMNP2] L. Ein, R. Lazarsfeld, M. Mustaţă, M. Nakamaye, and M. Popa, *Restricted volumes and base loci of linear series*, Amer. J. Math. **131** (2009), 607-651.

[F] W. Fulton, *Introduction to toric varieties*, Ann. of Math. Stud. **131**, Princeton Univ. Press, Princeton, NJ (1993).

[G] Y. Gongyo, *On the minimal model theory for dlt pairs of numerical log Kodaira dimension zero*, Math. Res. Lett. **18** (2011), 991-1000.

[J] S.-Y. Jow, *Okounkov bodies and restricted volumes along very general curves*, Adv. Math. **223** (2010), 1356-1371.

[KK] K. Kaveh and A. G. Khovanskii, *Newton convex bodies, semigroups of integral points, graded algebras and intersection theory*, Ann. of Math. (2) **176** (2012), 925-978.

[KL] A. Küronya and V. Lozovanu, *Local positivity of linear series on surfaces*, arXiv:1411.6205.

[La] R. Lazarsfeld,*Positivity in algebraic geometry I & II*, Ergeb. Math. Grenzgeb., **48** & **49** (2004), Springer-Verlag, Berlin.

[LM] R. Lazarsfeld and M. Mustaţă, *Convex bodies associated to linear series.* Ann. Sci. Ec. Norm. Super. (4) **42** (2009), 783-835.

[Le] B. Lehmann, *Comparing numerical dimensions*, Alg. and Num. Theory. **7** (2013), 1065-1100.

[N] N. Nakayama, *Zariski-decomposition and abundance*, MSJ Memoirs **14**. Mathematical Society of Japan, Tokyo, 2004.

[O1] A. Okounkov, *Brunn-Minkowski inequality for multiplicities*, Invent. Math. **125** (1996) 405-411.

[O2] A. Okounkov, *Why would multiplicities be log-concave?* in *The Orbit Method in Geometry and Physics*, Progr. Math. **213** (2003), Birkhuser Boston, Boston, MA, 329-347.

Department of Mathematics, Yonsei University, Seoul, Korea
E-mail address: `sungrakc@yonsei.ac.kr`

Samsung Advanced Institute of Technology, Suwon, Korea
E-mail address: `yoonsuk.hyun@samsung.com`

School of Mathematics, Korea Institute for Advanced Study, Seoul, Korea
E-mail address: `parkjh13@kias.re.kr`

Center for Geometry and Physics, Institute for Basic Science, Pohang, Korea
E-mail address: `leonwon@ibs.re.kr`